

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325339078>

Systematic review and meta-analysis of otitis media in Iran: Prevalence, etiology, antibiotic susceptibility, and complications

Article in *Indian Journal of Otology* · January 2018

DOI: 10.4103/indianjoto.IJOTOL_5_18z

CITATION

1

READS

36

6 authors, including:



Diana Sarokhani

Islamic Azad University Kermanshah Branch

31 PUBLICATIONS 116 CITATIONS

[SEE PROFILE](#)



Milad Azami

Ilam University of Medical Sciences

106 PUBLICATIONS 330 CITATIONS

[SEE PROFILE](#)



Mostafa Rezaei tavirani

Shahid Beheshti University of Medical Sciences

254 PUBLICATIONS 1,667 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Proteomic Analysis of Serum from Patients with Obsessive-Compulsive Disorder [View project](#)



Obstetrics, Gynecology and Infertility [View project](#)

Systematic Review and Meta-analysis of Otitis Media in Iran: Prevalence, Etiology, Antibiotic Susceptibility, and Complications

Safar-Ali Amiri-Andy, Diana Sarokhani, Milad Azami¹, Hossein Vazini², Mostafa Rezaei-Tavirani³, Ali Hasanpour Dehkordi⁴

Psychosocial Injuries Research Center, Ilam University of Medical Sciences, ¹Student Research Committee, Faculty of Medicine, Ilam University of Medical Sciences, ²Department of Nursing, Basic Sciences Faculty, Hamedan Branch, Islamic Azad University, Hamedan, ³Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, ⁴Department of Medical-Surgical, Faculty of Nursing and Midwifery, Shahrekord University of Medical Sciences, Shahrekord, Iran

Abstract

Otitis media (OM) is a relatively common infectious disease, and its chronic form may lead to complications such as hearing impairment. This study aims to determine the prevalence, etiology, antibiotic susceptibility, and complications of OM in Iran through meta-analysis. English language databases, including Science Direct, PubMed, Scopus, Web of Science, and Persian language databases, including SID, Magiran, Iranmedex, Medlib, as well as the Google Scholar search engine, were searched from 1994 to 2017, using the Mesh keywords: Otitis media, Middle ear, Causality, Etiology, and Iran. The heterogeneity assessment of the studies was done using Q-Cochran test and I² index. Considering the high heterogeneity of the studies, the random effects model was used to estimate the point prevalence with 95% confidence interval. Data were analyzed using STATA version 11.1 software. Fifty-one articles with a sample size of 10,675 were reviewed in this study, the most common types of OM involvement were right otitis (51% [95% confidence interval (CI): 33–68]), left otitis (44% [95% CI: 36–52]), and bilateral otitis (33% [95% CI: 7–59]). The most common bacterial etiologies involved in OM were *Pseudomonas aeruginosa* (26% [95% CI: 17–35]), *Proteus* (21% [95% CI: 0%–45%]), and *Staphylococcus* (20% [95% CI: 0–42]). The fungal etiology of OM was estimated to be 22% (95% CI: 4–40). The most common OM pathologies were ossicular changes (56% [95% CI: 27–85]), granulation (49% [95% CI: 18–80]), cholesteatoma (32% [95% CI: 13–51]), tympanosclerosis (17% [95% CI: 10–23]), and cholesterol granuloma (11% [95% CI: 0–26]). The results of this meta-analysis provide useful information on the epidemiology of OM in Iran to otorhinologists and health policymakers.

Keywords: Infection, Iran, meta-analysis, otitis media

INTRODUCTION

Otitis media (OM) is a general term for the expression of infections with various complications in the middle-ear region. OM are divided into three categories: acute OM, OM with effusion, and chronic suppurative OM (CSOM).^[1] CSOM account for a large number of patients referring to the ear, nose, and throat clinics and include a large number of surgical procedures.^[2,3]

In industrial countries, hearing loss (conductive and sensorineural) is the third-most common chronic illness after hypertension and arthropathy among the elderly, which has led to physical and mental problems for these people. Furthermore, according to our estimates, every year,

around 21,000 people (33/10 million people) die due to OM complications.^[4] Chronic illnesses lead to increased living costs, social problems, and impact on quality of life.^[5-9]

Chronic OM may cause ossicular erosion and as a result lead to major hearing loss.^[10] Hearing loss and otorrhea are common side effects of these diseases.^[2,3] Hearing loss occurs almost in all cases of chronic OM, which may reach 50–60 dB if

Address for correspondence: Dr. Mostafa Rezaei-Tavirani, Proteomics Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
E-mail: tavirany@yahoo.com

Access this article online

Quick Response Code:



Website:
www.indianjotol.org

DOI:
10.4103/indianjotol.INDIANJOTOL_5_18

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Amiri-Andy SA, Sarokhani D, Azami M, Vazini H, Rezaei-Tavirani M, Dehkordi AH. Systematic review and meta-analysis of otitis media in Iran: Prevalence, etiology, antibiotic susceptibility, and complications. *Indian J Otol* 2018;24:1-8.

ossicular erosion occurs.^[11] Considering the complications of OM, the recognition of causes and factors of OM is necessary.

The basis for the treatment of acute OM is the administration of antibiotics.^[12] Selective treatment of chronic OM is a surgical procedure and usually occurs after a common cold.^[13,14]

Different studies have suggested various pathologies as the causes of OM, most notably, granulation tissue, granulomatosis with polyangiitis (Wegener's), diabetes, vasculitis, cholesteatoma, cholesterol granuloma, tympanosclerosis, various viral infections, and ossicular changes.^[15-20] The ear cholesteatoma is referred to as epidermal inclusion cyst in the middle ear or mastoid. Due to the importance of the presence of cholesteatomas in the ear and the complications of its existence and failure to treat this problem, chronic inflammation of the middle ear is divided into two groups with cholesteatoma and without cholesteatoma.^[21] Due to the presence of creatine in a tissue space, cholesteatoma is prone to frequent infection, and *Pseudomonas aerogenosa* is its most common aerobic bacteria, while bacteroides species is the most common anaerobic bacteria.^[22]

Cholesterol granuloma as a foreign object causes granulation tissue.^[2] Granulation tissue is one of the most common pathologic findings in the ears in patients with chronic OM, which includes more than 2.3 cases of chronic OM without cholesteatoma.^[21] Tympanosclerosis is the formation of hyaline or calcified cartilage in the tympanic membrane and middle-ear cavity and is created secondary to inflammation of the middle ear or trauma, and histologically, hyaline deposition in the connective tissue is below the epithelium of the tympanic membrane and the middle ear.^[23]

Other chronic inflammatory pathologies of the middle ear are changes in the ossicular chain, which is often secondary to other pathologies.^[15,24] Bacteria and viruses also contribute to the development of OM. Common bacterial pathogens include *Pneumococcus*, *Haemophilus influenzae* and *Moraxella catarrhalis*.^[25-29] Of common viruses, rhinoviruses and the bacterial respiratory viruses can be mentioned, alone or as a pathogen.^[27-29]

Several researches have been conducted in Iran to study the etiology of OM, and the results are very different and, on the other hand, no systematic and meta-analytic reviews have been conducted on them. Therefore, performing a meta-analysis seems necessary. In the systematic review and meta-analysis, by combining different studies and increasing the number of studies and sample size involved in the analysis process, the 4 confidence intervals are reduced, and thus the results are more reliable.^[30] Therefore, the present study aims to investigate the OM etiology in Iran through systematic review and meta-analysis.

MATERIALS AND METHODS

Study protocol

The present study is a meta-analysis of the causes of OM in Iran. This study was conducted in accordance with the PRISMA protocol^[30] for systematic review and meta-analysis.

All the stages of this study were conducted independently by two researchers.

Search strategy

English language databases, including Science Direct, PubMed, Scopus, Web of Science, and Persian language databases, including SID, Magiran, Iranmedex, Medlib, as well as the Google Scholar search engine, were searched from 1994 to 2017, using the MeSH keywords: Otitis media, Middle ear, Causality, Etiology, and Iran. The combination of keywords was also performed using the AND/OR operator. The Persian equivalent of the keywords was also used for Persian language databases.

Inclusion and exclusion criteria

The main exclusion criterion of the study was the studies that examined the etiology of OM in Iran. Exclusion criteria were: (1) Irrelevance to the subject, (2) non-Iranian studies, (3) Studies with nonrandom sample size, (4) studies not conducted during the years 1994–2017, and (5) intervention articles, letters to editor, review articles, and case reports.

Qualitative assessment of studies

In order to assess the quality of studies, the standard and international STROBE checklist^[31] was used. This checklist contains 22 parts. Zero to two points were given to each part; therefore, the lowest and highest attainable points were 0 and 44, respectively. Studies that attained at least a score of 16 from the checklist entered the meta-analysis process.

Data extraction

To reduce reporting bias and data collection errors, two researchers independently performed data extraction and extracted data into a checklist that included the name of the first author, the study title, the sample size, the year and location of the study, the prevalence of each etiology involved in OM, the prevalence of complications of OM, and the sensitivity of OM.

Statistical analysis

In this study, the variances of each study were calculated using the binomial distribution formula. The heterogeneity evaluation was done by Q-Cochran test and I^2 index (I^2 index below 25% is low heterogeneity, between 25% and 75% is average heterogeneity, and 75% or above is high heterogeneity). Considering the high heterogeneity of the studies, a random effects model was used to estimate the point prevalence with 95% confidence interval (CI). All statistical analyses were performed using Streamline your internal and consumer-facing platforms with KitleyTech's industry leading custom software development services in Chicago. The significance level of the tests was considered to be $P < 0.05$.

RESULTS

Search results and properties of the studies that entered the meta-analysis process

In this systematic review and meta-analysis, 337 studies were identified in the initial search. After applying the inclusion

and exclusion criteria and qualitative assessment of the studies, 51 articles conducted from 1994 to 2017 entered the meta-analysis [Figure 1]. The properties of the studies are listed in Table 1.

In this study, the most common types of OM involvement were right otitis (51% [95% CI: 33–68]), left otitis (44% [95% CI: 36–52]), and bilateral otitis (33% [95% CI: 7–59]) [Table 2].

Otitis media etiology

The most common bacterial etiologies involved in OM were *Pseudomonas aeruginosa* (26% [CI 95%: 17–35]), *Proteus* (21% [95% CI: 0–45%]), and *Staphylococcus* (20% [95% CI: 0–42]). The fungal etiology of OM was estimated to be 22% (95% CI: 4–40).

Otitis media pathology

The most common OM pathologies were ossicular changes (56% [95% CI: 27–85]), granulation (49% [95% CI: 18–80]), cholesteatoma (32% [95% CI: 13–51]), tympanosclerosis (17% [95% CI: 10–23]), and cholesterol granuloma (11% [95% CI: 0–26]) [Table 2].

Otitis media complications

Among the complications of OM, the lowest prevalence was related to aural fullness (16% [95% CI: 9–23]) and dizziness (16% [95% CI: 5–28]), and the highest prevalence was associated with otorrhea (50% [95% CI: 0–1.32]) and moderate hearing loss (56% [95% CI: 49–62]) [Table 2].

Antibiotic susceptibility

In the assessment of antibiotic susceptibility of OM in Iran, the highest susceptibility was for gentamycin (86% [95% CI: 81–90]) and ciprofloxacin (81% [95% CI: 65–97]) and the lowest sensitivity was for penicillin (36% [95% CI: 14–59]) and amikacin (41% [95% CI: 0–88]) [Table 3].

Prevalence of surgical interventions

In the assessment of surgical interventions for chronic OM, the prevalence of high mastoidectomy was 47% (95% CI: 30–65), while the prevalence of low mastoidectomy and tympanoplasty was 24% (95% CI: 16–32) and 18% (95% CI: 14–23), respectively [Table 2].

DISCUSSION

The results of this meta-analysis on 51 articles with a sample size of 10,675 showed that the prevalence of right OM was more common than left otitis and bilateral otitis. The most common bacterial etiologies involved in OM were *P. aeruginosa* (26%), *Proteus* (21%), and *Staphylococcus* (20%), and the most common pathologies were ossicular changes (56%) and granulation (49%). Among the complications of OM, the lowest prevalence was associated with aural fullness (16%) and dizziness (16%) and the highest prevalence was associated with otorrhea (50%) and moderate hearing loss (56%). On investigating the antibiotic susceptibility for OM in Iran, the highest susceptibility was estimated for gentamycin (86%) and




Figure 1: Flowchart of the entry of studies into meta-analysis

Table 1: Data extracted from articles that entered the meta-analysis process

Author	Place	Year	Mean age	Method	n
Javadi M ^[15]	Tehran	2006	20-30	-	100
Fattahi Bafghi ^[32]	Tehran	2001-2003	15.26 (12.6)	Culturing	940
Sharifi Yazdi ^[33]	Tehran	2015	-	Culturing	102
Mohammadi ^[34]	Tabriz	1994-1998	-	-	400
Abtahi ^[35]	Esfahan	1996-1998	-	Culturing	154
Abshirini ^[36]	Ahvaz	2004	11-19	Culturing	100
Bakhshaei ^[37]	Mashhad	2004-2005	<6	Culturing	1161
Maymane Jahromi ^[38]	Mashhad	2004-2009	23.08 (11.35)	-	70
Khorshidi ^[39]	Kashan	2001-2002	-	Culturing	50
Malakooti ^[40]	Tehran	1997	-	-	126
Pirzade ^[41]	Ardebil	1999-2001	-	Culturing	60
Toutounchi ^[42]	Tabriz	1998-2000	27	-	37
Mozafari ^[43]	Zanjan	2004-2005	18-45	Culturing	100
Sadooghi ^[44]	Tehran	2001	-	-	212
Rajati Haghi ^[45]	Mashhad	2007-2008	9-70	-	97
Naeimi ^[46]	Mashhad	2007-2008	28.4	Culturing	62
Nahaei ^[47]	Tabriz	1998	-	-	50
Iranfar ^[48]	Esfahan	2002-2003	-	-	90
Soltan Dallal ^[49]	Tehran	2013	-	Culturing	102
Soltan Dallal ^[50]	Tehran	2013	-	Culturing	102
Asadi ^[51]	Tehran	2013-2014	2-26	PCR	60
Nader Poor ^[52]	Tabriz	2001-2007	30.8	-	166
Aynol Ghazati ^[53]	Mashhad	2001-2003	-	-	116
Khoshdel ^[54]	Shahrekord	2007	<5	Culturing	224
Sabory ^[55]	Kermanshah	2012	-	Culturing	554
Delsouz Bahri ^[56]	Tehran	2013-2014	-	Culturing	50
Delsouz Bahri ^[56]	Tehran	2013-2014	-	Culturing	50
Ghiassi S ^[10]	Tehran	2008-2009	-	PCR	196
Kafshdar Jalali ^[57]	Lahijan	2009-2010	20-36	-	120
Jolodarian ^[58]	Ahvaz	2002-2006	22.87	Culturing	548
Dadgarnia ^[59]	Yazd	1998-2001	-	-	80
Namaei ^[60]	Birjand	2012-2014	-	-	100
Yeganeh Moghadam ^[61]	Kashan	2010-2012	6.6	Culturing	90
Karimi ^[62]	Yazd	1995-1996	-	Culturing	120
Ebrahimi ^[63]	Tehran	2015	-	Culturing	60
Ebrahimi ^[63]	Tehran	2015	-	PCR	60
Aseiri ^[64]		1999	-	-	775
Farhadi ^[65]	Tehran	2000	-	Culturing	52
Mirvakili ^[66]	Yazd	2004-2006	32.4	-	120
Ghaznavi Rad ^[67]	Arak	2003	-	Culturing	42
Jalali ^[15]	Rasht	2004-2005	1-14	Culturing	74
Gharibpour ^[68]	Tehran	2010-2011	-	Culturing	50
Gharibpour ^[68]	Tehran	2010-2011	-	PCR	50
Salehe ^[69]	Ahvaz	2011	-	-	76
Saki ^[70]	Ahvaz	1996	-	-	2000
Nourizadeh ^[71]	Mashhad	2015	-	-	30
Khoramrooz ^[72]	Tehran	2009-2010	7.2	-	50
Farajzadah Sheikh ^[73]	Ahvaz	2014	1-15	Culturing	45
Farajzadah Sheikh ^[73]	Ahvaz	2014	1-15	PCR	45
Abshirini ^[74]	Ahvaz	2012	-	-	30
Emaneini ^[75]	Tehran	2009-2010	-	-	45
Khoramrooz ^[76]	Tehran	2009-2010	7.01	Culturing	63
Khoramrooz ^[76]	Tehran	2009-2010	7.01	PCR	63

Contd...

Table 1: Contd...

Author	Place	Year	Mean age	Method	n
Jalali ^[77]	Rasht	2005–2006	1-14	Culturing	62
Ettehad ^[78]	Ardebil	2003–2004	21-30	Culturing	61
Mozafari Nia ^[79]	Kerman	2011	-	-	117
Faramarzi ^[80]	Tehran	2000–2006	29.7	-	116

PCR: Polymerase chain reaction

Table 2: Results of the meta-analysis of articles: Causes of otitis media in Iran

Subgroups	Number of studies	Prevalence	Low	Up	P	I ² (%)
The prevalence of OM						
Bilateral otitis	5	33	7	59	0.000	96.2
Left otitis	2	44	36	52	0.307	4.0
Right otitis	2	51	33	68	0.052	73.4
The prevalence of OM etiology						
<i>Pseudomonas aeruginosa</i>	14	26	17	35	0.000	96.5
<i>Staphylococcus aureus</i>	12	18	12	23	0.000	87.9
<i>Staphylococcus</i>	3	20	0	42	0.000	97.6
Golden <i>Staphylococcus</i>	2	18	0	43	0.000	93.6
<i>Staphylococcus epidermidis</i>	3	9	1	18	0.003	83.1
<i>Haemophilus influenzae</i>	12	13	7	18	0.000	89.5
<i>Proteus</i>	9	21	0	45	0.000	99.6
<i>aspergillus</i>	2	9	3	15	0.01	84.9
<i>Klebsiella pneumoniae</i>	6	5	2	7	0.005	70.4
<i>Streptococcus pneumoniae</i>	9	11	9	14	0.001	68.7
<i>Pneumococcus</i>	9	8	4	13	0.000	86.1
<i>Moraxella catarrhalis</i>	9	6	3	9	0.007	62.0
<i>Branhamella catarrhalis</i>	2	14	9	18	0.801	0.0
<i>Candida albicans</i>	2	6	2	11	0.982	0.0
<i>Escherichia coli</i>	6	9	6	12	0.602	0.0
<i>Alloioicoccus otitidis</i>	11	25	15	35	0.000	93.5
Fungus	6	22	4	40	0.000	98.1
Cholesteatoma	15	32	13	51	0.000	99.4
Cholesterol granuloma	2	11	0	26	0.020	81.6
Granulation	5	49	18	80	0.000	99.0
Tympanosclerosis	6	17	10	23	0.000	78.7
Ossicular changes	4	56	27	85	0.000	98.2
The prevalence of OM complications						
Otorrhea	2	50	0	1.32	0.000	99.8
Aural fullness	1	16	9	23	-	100
Dizziness	4	16	5	28	0.000	92.0
Tinnitus	4	29	4	54	0.000	97.7
Hearing loss	4	31	3	65	0.000	99.6
Mild hearing loss	2	22	17	28	0.599	0.0
Moderate hearing loss	2	56	49	62	0.658	0.0
Severe hearing loss	3	37	0	76	0.000	98.9
The prevalence of surgical interventions						
Tympanoplasty	3	18	14	23	0.329	10.0
High mastoidectomy	3	47	30	65	0.000	91.0
Low mastoidectomy	1	24	16	32	-	100

ciprofloxacin (81%) antibiotics and the lowest susceptibility was for penicillin (36%) and amikacin (41%) antibiotics.

OM is considered an indigenous disease in the northern region of Australia and, in the villages of this area, all

children develop the infection several days after birth, and the pathogens of *Streptococcus pneumoniae*, *H. influenzae*, and *M. catarrhalis* are observed in them.^[81] In a study in Israel, the prevalence of chronic OM was 95% and 41% of those with chronic OM had cholesteatoma.^[82] In another study

Table 3: Antibiotic susceptibility administered for patients with otitis media

Drugs	Number of studies	Prevalence	Low	Up	P	I ² (%)
Susceptibility to drug						
Azithromycin	6	53	40	65	0.000	96.2
Co-trimoxazole	4	44	18	70	0.000	98.0
Cefixime	5	48	28	67	0.000	97.3
Amikacin	3	41	0	88	0.000	99.0
Chloramphenicol	2	52	28	76	0.002	89.5
Tetracycline	2	65	49	81	0.015	83.1
Carbenicillin	2	48	6	91	0.000	97.8
Streptomycin	2	49	31	66	0.027	79.6
Vancomycin	3	63	40	87	0.000	95.3
Ceftriaxone	3	48	25	72	0.000	96.5
Penicillin	4	36	14	59	0.000	98.7
Amoxicillin	4	54	27	81	0.000	99.3
Gentamicin	2	86	81	90	0.839	0.0
Ciprofloxacin	3	81	65	97	0.000	93.0

by Sommerfleck in Argentina in 2012, 324 patients were diagnosed with acute OM. The most significant pathogens in patients were *S. pneumoniae* (39.5%), *H. influenza* (37.4%), *M. catarrhalis* (6.1%), and *S. pneumoniae* (3%).^[83] In a study in Malaysia in 1999, which was conducted among 382 patients, the most common organisms were *P. aeruginosa* (27.2%) and *Staphylococcus aureus* (23.6%).^[84]

In Japan, golden *Staphylococcus* has grown more and more significantly over 16 years in Chronic suppurative otitis media (CSOM) patients, compared with other microorganisms.^[85] A study by Nyembu *et al.* in Congo in 2003 on ear effusions of children showed that *Proteus mirabilis* (23%), *P. aeruginosa* (22%), and *Citrobacter koseri* (20%) had the highest prevalence of microorganism in CSOM children.^[86] In a study by Moshi *et al.* in Tanzania in 2000, by investigating 176 effusions from 150 patients with CSOM, *P. aeruginosa* (51%), *S. aureus* (17.2%), *P. mirabilis* (13.2%), *Klebsiella pneumoniae* (8%), and *Escherichia coli* (5.8%) were isolated from culture media.^[87] Sharma *et al.* in Nepal investigated 322 effusions from 250 patients with CSOM and showed that *P. aeruginosa* (36.4%) and *S. aureus* (30.2%) had the highest prevalence of microorganisms in patients with CSOM.^[88]

One of the limitations of this study was the limited statistical population of Iran, and some studies only focused on the prevalence of OM.

CONCLUSION

According to the results of this study, continuing education for community members and stakeholders in relation to OM, antibiotic susceptibility to effective treatment, prevention of complications, antibiotic resistance, and reduction of treatment costs are essential.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Gates GA, Klein JO, Lim DJ, Mogi G, Ogra PL, Pararella MM, *et al.* Recent advances in otitis media 1. Definitions, terminology, and classification of otitis media. *Ann Otol Rhinol Laryngol Suppl* 2002;188:8-18.
- Chole R, Sudhoff H. Chronic otitis media, mastoiditis, and petrositis. In: Cummings CW, Flint PW, Haughey BH, Robbins KT, Thomas JR, Harker LA, *et al.*, editors. *Cummings Otolaryngology Head and Neck Surgery*. 4th ed. Philadelphia: Elsevier; 2005. p. 2988-3012.
- Wang HM, Lin JC, Lee KW, Tai CF, Wang LF, Chang HM, *et al.* Analysis of mastoid findings at surgery to treat middle ear cholesteatoma. *Arch Otolaryngol Head Neck Surg* 2006;132:1307-10.
- Monasta L, Ronfani L, Marchetti F, Montico M, Vecchi Brumatti L, Bavcar A, *et al.* Burden of disease caused by otitis media: Systematic review and global estimates. *PLoS One* 2012;7:e36226.
- Hasanpour Dehkordi A. Influence of yoga and aerobics exercise on fatigue, pain and psychosocial status in patients with multiple sclerosis: A randomized trial. *J Sports Med Phys Fitness* 2016;56:1417-22.
- Hasanpour-Dehkordi A, Khaledi-Far A, Khaledi-Far B, Salehi-Tali S. The effect of family training and support on the quality of life and cost of hospital readmissions in congestive heart failure patients in Iran. *Appl Nurs Res* 2016;31:165-9.
- HassanpourDehkordi A, Mohammadi N, NikbakhatNasrabadi A. Hepatitis-related stigma in chronic patients: A qualitative study. *Appl Nurs Res* 2016;29:206-10.
- Mohammadi N, HassanpourDehkordi A, NikbakhatNasrabadi A. Iranian patients with chronic hepatitis struggle to do self-care. *Life Sci J* 2013;10:457-62.
- Ghiasi B, Sarokhani D, Dehkordi AH, Sayehmiri K, Heidari MH. Quality of Life of patients with chronic kidney disease in Iran: Systematic Review and Meta-analysis. *Indian Journal of Palliative Care*. 2018;24:104.
- Ghiasi S, Seyed Toutounchi J. Prevalence of ossicular lesions in chronic otitis media. *Med J Tabriz Univ Med Sci Health Serv* 2014;36:66-9.
- Ibrahim SI, Cheang PP, Nunez DA. Incidence of meningitis secondary to suppurative otitis media in adults. *J Laryngol Otol* 2010;124:1158-61.
- Block SL, Schmier JK, Notario GF, Akinlade BK, Busman TA, Mackinnon GE 3rd, *et al.* Efficacy, tolerability, and parent reported outcomes for cefdinir vs. high-dose amoxicillin/clavulanate oral suspension for acute otitis media in young children. *Curr Med Res Opin* 2006;22:1839-47.
- Arroll B. Common cold. *BMJ Clin Evid* 2008. pii: 1510.

14. Yingxi L, Sheng L, Xiuzhen S. Numerical analysis of ossicular chain lesion of human ear. *J Acta Mech Sin* 2008;25:241-7.
15. Javadi M, Mohebi S. Pathologic findings in 100 patients with chronic otitis media in Hazrat-e-Rasool Akram Hospital. *J Iran Univ Med Sci* 2007;14:81-7.
16. Amiri M. Diabetes mellitus type 2; an international challenge. *Ann Res Dial*. 2016;1:e04.
17. Dey B, Dange P, Ganesh RN, Parameswaran S, Pillai PP. Immune-complex deposits in anti-neutrophil cytoplasmic antibody associated crescentic glomerulonephritis; a report of two cases. *Immunopathologia Persa*, 2017;3:e09, 10.15171/ipp.2017.01.
18. Kronbichler A. Boon and bane of remission induction with rituximab in ANCA-associated vasculitis: Lessons learned from the RAVE-ITN follow-up study. *J Renal Inj Prev* 2014;3:9-10.
19. Mortazavi M, Nasri H. Granulomatosis with polyangiitis (Wegener's) presenting as the right ventricular masses: A case report and review of the literature. *J Nephropathol* 2012;1:49-56.
20. Rodríguez-Morales AJ, Sánchez-Duque JA. Preparing for next arboviral epidemics in Latin America; who can it be now? – Mayaro, Oropouche, West Nile or Venezuelan equine encephalitis viruses. *J Prev Epidemiol* 2018;3:e01.
21. da Costa SS, Paparella MM, Schachern PA, Yoon TH, Kimberley BP. Temporal bone histopathology in chronically infected ears with intact and perforated tympanic membranes. *Laryngoscope* 1992;102:1229-36.
22. Chole RA, Faddis BT. Evidence for microbial biofilms in cholesteatomas. *Arch Otolaryngol Head Neck Surg* 2002;128:1129-33.
23. Dew L, Shelton C. Complication of temporal bone infection. In: Cummings CW, Flint PW, Harker LA, Haughey BH, Richardson MA, Robinson KT, *et al.*, editors. *Otolaryngology Head and Neck Surgery*. 3rd ed. St. Louis: Mosby; 1998. p. 3013-38.
24. Shahreza FD. From oxidative stress to endothelial cell dysfunction. *J Prev Epidemiol* 2016;1:e04.
25. Couloigner V, Levy C, François M, Bidet P, Hausdorff WP, Pascal T, *et al.* Pathogens implicated in acute otitis media failures after 7-valent pneumococcal conjugate vaccine implementation in France: Distribution, serotypes, and resistance levels. *Pediatr Infect Dis J* 2012;31:154-8.
26. Wen R, Deng Q, Sun C, Gao S, Tao J, Luo R, *et al.* Pathogenic bacteria distribution and drug susceptibility in children with acute otitis media in pearl river Delta. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi* 2011;25:884-7.
27. Casey JR, Pichichero ME. Changes in frequency and pathogens causing acute otitis media in 1995-2003. *Pediatr Infect Dis J* 2004;23:824-8.
28. Faden H. The microbiologic and immunologic basis for recurrent otitis media in children. *Eur J Pediatr* 2001;160:407-13.
29. Pumarola F, Marès J, Losada I, Minguella I, Moraga F, Tarragó D, *et al.* Microbiology of bacteria causing recurrent acute otitis media (AOM) and AOM treatment failure in young children in Spain: Shifting pathogens in the post-pneumococcal conjugate vaccination era. *Int J Pediatr Otorhinolaryngol* 2013;77:1231-6.
30. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
31. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, *et al.* Strengthening of reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *BMJ* 2007;335:806-8.
32. Malakooti B, Mahdizadeh J, Amani Gashnigani S, Fathami S. Effects of eustachian tube function in outcome of graft in tympanoplasty or tympanomastoidectomy for chronic otitis media. *Tehran Univ Med J* 2000;58:11-3.
33. Pirzadeh A, Ettehad G. A study of etiologies of chronic otitis media at patients referring to ear, throat and nose clinics of Ardabil University of Medical Sciences, 2000-2001. *J Ardabil Univ Med Sci* 2002;2:10-3.
34. Toutounchi SJ, Abbaszadeh P, Barzgar M. Degree of hearing loss and different pathology of middle ear. *J Ardabil Univ Med Sci* 2005;5:119-23.
35. Mozafari N, Talei S, Amirmoghaddami H, Talei S. Culture and antibiogram of aerobic bacteria causing chronic otitis media in Zanjan. *ZUMS J* 2006;14:52-9.
36. Sadooghi M, Soofizade M. Evaluation of chronic otitis media in patients referred to Amir Alam hospital during the first six month of 2001. An analysis of 212 cases. *J Arc* 2003;16:39-42.
37. Rajati Haghi M, Ghasemi M, Bakhshae M, Taghati A, Shahabi Poor A. Audiometry and ossicles in patients with chronic otitis media. *Iran J Otorhinolaryngol* 2009;21:73-7.
38. Naeimi M, Ghazvini K, Shakeri M, Kaboli M, Bagheri M. Study of bacteremia incidence after middle ear surgery. *Iranian journal of ears, throat, nose and larynx. Iran J Otorhinolaryngol* 2008;20:177-82.
39. Nahaei M, Kazem Nejad M. The prevalence of various lesions in the middle ear and their relationship with preoperative ODM results in 50 surgeries performed on middle ear of adolescents and adults admitted to Imam Khomeini hospital in Tabriz in 1998. *Med J Tabriz Univ Med Sci* 2000;34:63-70.
40. Iranfar K, Ragha M, Abtahi S. The effect of middle ear surgery on sensorineural hearing in patients with chronic otitis media. *J Kermanshah Univ Med Sci* 2004;8:22-8.
41. Soltan Dallal M, Rahimi Forushani A, Heidarzadeh S, Jabbari H, Sedigh Maroufi S, Sharifi Yazdi M. Identification and determination antimicrobial resistance pattern of *Haemophilus influenza* isolated from patients with otitis media. *RJMS* 2014;21:17-24.
42. Soltan Dallal M, Jabari H, Rahimi Forushani A, Heidarzadeh S, Afrogh P, Sharifi Yazdi M. Frequency and resistance patterns of *Streptococcus pneumoniae* in acute otitis media. *J Mazandaran Univ Med Sci* 2013;23:28-35.
43. Asadi A, Gudarzi M, Gudarzi H, Houry H, Ebrahimi N, Delsoz Bahri T. Investigating the frequency and resistance to treatment of *Streptococcus pneumoniae* in patients with otitis. *Koomesh* 2016;17:814-20.
44. Nader Poor M, Musavi Aghdas M. Cholesteatoma surgery results in 166 patients. *Med J Tabriz Univ Med Sci* 2010;22:73-7.
45. Aynol Ghazati S, Sharifi Dalooei M, Adami Dehkordi M, Nikzade M. Examining secretory chronic otitis media in teaching hospitals of Islamic Azad University, Mashhad, 2001-2003. *Med J Islamic Azad Univ Mashhad* 2005;3:36-42.
46. Khoshdel A, Imani R, Saedi A, Kheire S, Hamidi M, Kasiri K, *et al.* The prevalence of *Streptococcus pneumonia* and its penicillin resistance pattern in children less than five years old from Shahrekord Iran 2007. *J Shahrekord Univ Med Sci* 2009;10:89-95.
47. Sabory T, Ghadiri K, Abiri R, Elahi A, Poormohammadi S, Gharib A. Incidence of nasopharyngeal carriers of *Streptococcus pneumoniae* and antibiotic resistance in the children in Kermanshah 2012. *J Nurs Educ* 2016;4:90-7.
48. Delsouz Bahri T, Goudarzi M, Dabiri Satri S, Ebrahimi N, Asadi A, Ghafoori S, *et al.* Comparison of culture and multiplex PCR in detection of fastidious bacteria associated with otitis media among suspected patient admitted to Amir-Alam Hospital. *Res Med* 2017;40:192-6.
49. Kafshdar Jalali A, Salamat Zade A. Examining bacterial causes of chronic otitis media and their antibiotic susceptibility in Lahijan hospitals. *Tabriz University of Medical Sciences and Health Services First International Congress and 4th Conference on Health and Education*; 2011.
50. Jolodarian P, Abshirini H. Evaluation of Epidemiological Findings of Cholesteatomas in Patients with Chronic Otitis Media Admitted to Going Under Surgery in the Ear and Nose Centers of Imam Khomeini Hospital in Ahvaz between 2002 and 2006. *Jundishapur University of Medical Sciences & Health Services*; 2009.
51. Asadi A, Gudarzi M, Gudarzi H, Houry H, Ebrahimi N, Delsoz Bahri T. Investigating the frequency and resistance to treatment of *Streptococcus pneumoniae* in patients with otitis. *Koomesh* 2016;17:814-20.
52. Namaei M, Mehramiz M, Ghannadkafi M, Mofatteh MR. Prevalence of bacterial causes of chronic otitis media with effusion and their drug resistance in Birjand. *J Birjand Univ Med Sci* 2015;22:59-66.
53. Yeganeh Moghadam A, Namazi M, Moravveji S, Khorshidi A, Abbasi Moghadam A. Bacteriological assessment of resistant effusional otitis media discharge among the 2-16 year-old children admitted to Matini Hospital in Kashan, 2011-2013. *Feyz* 2014;18:469-76.
54. Karimi G, Moradi A. Study of microbial effusion of the ear in chronic otitis media and its prevalence among 120 hospitalized patients and outpatients. *J Shahid Sadoughi Univ Med Sci Yazd* 2001;9:53-6.
55. Ebrahimi N, Goudarzi H, Goudarzi M, Asadi A, Delsouz Bahri T. Frequency of alloiococcus otitis in patients with otitis media by culture

- and PCR method. *Res Med* 2015;39:36-40.
56. Asiri S, Hasham A, al Anazy F, Zakzouk S, Banjar A. Tympanosclerosis: Review of literature and incidence among patients with middle-ear infection. *J Laryngol Otol* 1999;113:1076-80.
57. Farhadi M, Tabatabai A, Shamshiri A, Maleki A, Masjedani F, Bojary Nassrabadi M, *et al.* Comparative study of Aerobic and anaerobic bacteria and fungi involved in sinusitis and otitis media cases from patients under operated in ENT ward at Rasool Hospital. *Razi J Med Sci* 2000;7:44-9.
58. Mirvakili A, Baradaranfar M, Hasani A, Jafari R. Sensorineural hearing loss in chronic otitis media. *JSSU* 2008;15:21-8.
59. Ghaznavi Rad E, Zareie R, Jafari A, Palizvan MR, Jourabchi A, Moini L, *et al.* Frequency and risk factors of *Moraxella catarrhalis* infection in patients at Vali-Asr and Amirkabir hospitals since August 2003 to March 2004. *Yafte* 2006;7:117-22.
60. Gharibpour F, Khoramrooz S, Mirsalehian A, Emaneini M, Jabalameli F, Darban-Sarokhalil D, *et al.* Isolation and detection of *Alloicoccus otitidis* in children with otitis media with effusion using culture and PCR methods. *J Mazand Univ Med Sci* 2013;23:52-60.
61. Salehe F, Abshirini H. To Evaluate the Relationship Between Auditory Findings and Status of Ossicular Chain During Surgery in Patients with Chronic Otitis Media. These is of Jundishapur University of Medical Sciences and Health Services, Ahvaz; 2012.
62. Saki N, Nik Akhlagh S, Modarres Dezfali S. Determination of the prevalence of otitis media with effusion in the first grade elementary school students of Ahvaz in the winter of 1996. *J Med Sci* 2005;4:211-5.
63. Nourizadeh N, Ghazvini K, Gharavi V, Nourizadeh N, Movahed R. Evaluation of nasopharyngeal microbial flora and antibiogram and its relation to otitis media with effusion. *Eur Arch Otorhinolaryngol* 2016;273:859-63.
64. Khoramrooz SS, Mirsalehian A, Imaneini H, Jabalameli F, Sharifi A, Aligholi M, *et al.* Characterization of *Alloicoccus otitidis* strains isolated from children with otitis media with effusion by pulsed-field gel electrophoresis. *Int J Pediatr Otorhinolaryngol* 2012;76:1658-60.
65. Farajzadah Sheikh A, Saki N, Roointan M, Ranjbar R, Yadyad MJ, Kaydani A, *et al.* Identification of *Alloicoccus otitidis*, *Streptococcus pneumoniae*, *Moraxella catarrhalis* and *Haemophilus influenzae* in children with otitis media with effusion. *Jundishapur J Microbiol* 2015;8:e17985.
66. Abshirini H, Khosravi A, Ghazipour A, Yavari M, Hashemzade M, Salehi F. Evaluation of cholesteatoma microbiologic finding in patients with chronic. *Health Med* 2012;6:1684-7.
67. Emaneini M, Gharibpour F, Khoramrooz SS, Mirsalehian A, Jabalameli F, Darban-Sarokhalil D, *et al.* Genetic similarity between adenoid tissue and middle ear fluid isolates of *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis* from Iranian children with otitis media with effusion. *Int J Pediatr Otorhinolaryngol* 2013;77:1841-5.
68. Khoramrooz SS, Mirsalehian A, Emaneini M, Jabalameli F, Aligholi M, Saedi B, *et al.* Frequency of *Alloicoccus otitidis*, *Streptococcus pneumoniae*, *Moraxella catarrhalis* and *Haemophilus influenzae* in children with otitis media with effusion (OME) in Iranian patients. *Auris Nasus Larynx* 2012;39:369-73.
69. Jalali M, Rezaie A, Kousha A, Saadat F, Banan R. Detection of fungal DNA in the middle ear effusion of patients suffering from otitis media with effusion. *Iran J Public Health* 2008;37:109-13.
70. Ettihad G, Refahi S, Nemmati A, Pirzadeh A, Daryani A. Microbial and antimicrobial susceptibility patterns from patients with chronic otitis media in Ardebil. *Int J Trop Med* 2006;1:62-5.
71. Mozafari Nia K, Sepehri G, Khatmi H, Shakibaie MR. Isolation and antimicrobial susceptibility of bacteria from chronic suppurative otitis media patients in Kerman, Iran. *Iran Red Crescent Med J* 2011;13:891-4.
72. Faramarzi A, Motasaddi-Zarandy M, Khorsandi MT. Intraoperative findings in revision chronic otitis media surgery. *Arch Iran Med* 2008;11:196-9.
73. Gibney KB, Morris PS, Carapetis JR, Skull SA, Smith-Vaughan HC, Stubbs E, *et al.* The clinical course of acute otitis media in high-risk Australian Aboriginal children: A longitudinal study. *BMC Pediatr* 2005;5:16.
74. Fliss DM, Shoham I, Leiberman A, Dagan R. Chronic suppurative otitis media without cholesteatoma in children in Southern Israel: Incidence and risk factors. *Pediatr Infect Dis J* 1991;10:895-9.
75. Pellegrini S, Gonzalez Macchi ME, Sommerfleck PA, Bernaldez PC. Intratemporal complications from acute otitis media in children: 17 cases in two years. *Acta Otorrinolaringol Esp* 2012;63:21-5.
76. Indudharan R, Haq JA, Aiyar S. Antibiotics in chronic suppurative otitis media: A bacteriologic study. *Ann Otol Rhinol Laryngol* 1999;108:440-5.
77. Nakagawa T, Yadhisa O, Komunes S. A 19-year survey of changes in bacterial isolate associated with chronic suppurative otitis media. *Ear Arch Otorhinolaryngol* 1994;1:27-32.
78. Nyembue DT, Tshiswaka JM, Sabue MJ, Muyunga CK. Bacteriology of chronic suppurative otitis media in Congolese children. *Acta Otorhinolaryngol Belg* 2003;57:205-8.
79. Moshi NH, Minja BM, Ole-Lengine L, Mwakagile DS. Bacteriology of chronic otitis media in Dar es Salaam, Tanzania. *East Afr Med J* 2000;77:20-2.
80. Sharma S, Rehan HS, Goyal A, Jha AK, Upadhyaya S, Mishra SC, *et al.* Bacteriological profile in chronic suppurative otitis media in Eastern Nepal. *Trop Doct* 2004;34:102-4.
81. Fattahi Bafghi A, Paeyvandi A. The study of pathogens in the secretion of chronic infectious diseases in the middle ear. *Sci J Med Organ Islam Repub Iran* 2010;28:397-402.
82. Sharifi Yazdi M, Heidarzade S, Vahedi S, Rahimi Forushani A, Jabari H, Soltan Dallal M. Identification and determination antimicrobial resistance pattern of *Staphylococcus aureus* and *Pseudomonas aeruginosa* isolated from patients with otitis media. *J Islam Repub Iran Med Counc* 2015;33:91-100.
83. Mohammadi GA, Sayah Melli M. The efficacy of tympanomastoid surgery for prevention of infection in active chronic otitis media. *Med J Tabriz Univ Med Sci Health Serv* 2002;55:55-9.
84. Abtahi S, Hashemi S. Bacteriological study of middle ear discharge in secretory otitis media. *Iran J Otorhinolaryngol* 2002;14:30-3.
85. Abshirini H. Culturing and antibiogram of aerobic bacteria causing chronic otitis media. *Iran J Otorhinolaryngol* 2004;16:28-37.
86. Nyembu D, Tshiswaka J, Sabue M, Mu yunga C. Bacteriology of chronic suppurative otitis media in Congolese children. *Acta otorhinolaryngol Belg* 2003;57:205-9.
87. Maymane Jahromi A, Nezhad Aryani FH, Arab Khani R, Nezhad Aryani SH. Investigating the frequency of cholesteatoma in patients with chronic otitis media referring to the eardrum, throat and nose center of imam reza hospital of Mashhad during 2004-2009. *Iran J Otorhinolaryngol* 2009;21:21-4.
88. Sharma S, Rehan H, Goyla A, Jha A, Upadhyaya S, Mishra S. Bacteriological profile in chronic suppurative otitis media in Eastern Nepal. *Trop Doct* 2004;34:102-4.